

# The Impact of Social Distancing for Severe Acute Respiratory Syndrome Coronavirus 2 on Respiratory Syncytial Virus and Influenza Burden

Kathryn M. Edwards<sup>®</sup>

Department of Pediatrics, Division of Pediatric Infectious Diseases, Vanderbilt University Medical Center, Nashville, Tennessee, USA

(See the Major Article by Friedrich et al on pages 2071–5, Brief Reports by Nolen et al on pages 2196–8 and Yeoh et al on pages 2199–202 and the Editorial Commentary by Covin and Rutherford on pages e993–4.)

**Keywords.** social distancing; influenza; respiratory syncytial virus; SARS Cov2.

Each year pediatricians dread the annual arrival of respiratory syncytial virus (RSV) and influenza virus because they are both associated with increased respiratory illnesses and resultant hospitalizations. RSV is the major cause of pneumonia in young children and is responsible for yearly global morbidity and mortality [1, 2]. Influenza virus also has a major impact, with >100 pediatric deaths each year attributable to influenza in the United States, and the global burden is even more impactful [3, 4]. Thus, 3 articles appearing in this issue of *Clinical Infectious Diseases* demonstrate impressive declines in both RSV and influenza associated with mitigation strategies implemented in response to the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic. They occur in 3 disparate regions of the world but provide some common perspectives [5–7].

Infants in the remote Yukon-Kuskokwim Delta (YKD) region of Alaska

have the highest rates of RSV hospitalization in the United States, reaching 259 per 1000 infants during the winter season [8]. Since infants in the YKD region are all hospitalized in 1 hospital and since comprehensive viral surveillance for RSV has been conducted for >25 years, the impact of mitigation strategies on RSV burden can be comprehensively assessed [9]. In late February 2020, infants with RSV were hospitalized in the region, but there were no SARS-CoV-2 isolates detected in the community. However, on 3 March social distancing mandates were implemented in the region: school closure, travel limitations to the region, and the closure of nonessential businesses and social distancing from nonhousehold members. The rate of RSV hospitalizations in the region precipitously declined with no RSV hospitalizations occurring after the second week in April. Remarkably, among the 25 previous RSV seasons, such an occurrence had never been seen. Overall hospitalizations for other illnesses had not changed and no children were admitted with severe respiratory illness that had resulted from avoidance of care. These observations suggested that social distancing and travel restrictions markedly reduced the contagion.

The second observations are reported from Brazil and also assessed the impact of mitigation strategies on acute

bronchiolitis, largely secondary to RSV. In Brazil, acute bronchiolitis is responsible for 6% of the total hospitalizations in children <1 year of age, with a peak incidence between the months of February and August [10]. Mitigation measures to prevent SARS-CoV-2 infection were implemented in Brazil in the middle of March and included social distancing, restrictions on business and nonessential services, and closure of schools. Data regarding acute bronchiolitis hospitalizations in infants <1 year of age were compared for the past 4 years. A significant reduction in the number of hospitalizations for acute bronchiolitis in infants aged <1 year was noted, with a >70% decline when comparing the rates from earlier years. These data also highlight the impact of social distancing and business and school closures on reducing the transmission of RSV in young children.

The final manuscript is from Western Australia, where mitigation strategies to prevent coronavirus disease 2019 (COVID-19) were implemented at the end of March, just prior to the Southern Hemisphere winter, the usual period of peak local RSV and influenza infections [11]. Since the RSV and influenza samples from all public hospitals and emergency departments in the region have been tested at a single laboratory since 2012, data on numbers of isolates

Received 2 October 2020; editorial decision 4 October 2020; published online 11 October 2020.

Correspondence: Correspondence: K. M. Edwards, Department of Pediatrics, D 7227 Medical Center North, Vanderbilt University Medical School, Nashville, TN 37232 (kathryn.edwards@vumc.org).

**Clinical Infectious Diseases**® 2021;72(12):2076–8

© The Author(s) 2020. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com. DOI: 10.1093/cid/ciaa1543

detected could be compared over time. The mitigation strategies included border closure to nonresidents in March and stay-at-home restrictions with school closure from the end of March until the end of April. Schools were then reopened with no restrictions on attendance. In earlier years, the annual number of RSV cases ranged from 532 to 707 and the annual number of influenza cases ranged from 134 to 498. For the first 13 weeks of 2020, prior to the local mitigation restrictions, RSV and influenza detections were comparable to earlier seasons. However, after mitigation strategies were implemented, influenza and RSV quickly declined and remained low, even in the face of school reopening. When compared with previous prepandemic years, RSV detections were 98% lower and influenza detections 99% lower than in earlier years. This seems even more remarkable since the decline persisted when school resumed. It is widely known that school-aged children are the major transmission vectors to younger children for both influenza and RSV [12, 13]. Although social distancing and attention to hygiene are likely contributing factors, the marked reduction in the number of RSV and influenza cases is consistent with reduced transmission from travel restrictions and closure of external borders. Reductions in Southern Hemisphere influenza detections recently reported from both the World Health Organization and the Centers for Disease Control and Prevention surveillance data reflect these same trends [14, 15].

Although these reports focused on RSV and influenza, additional studies demonstrated that other respiratory illnesses have also decreased with social mitigation strategies. An impressive report from Paris demonstrated the impact of school closures and social distancing on the reduction in pediatric hospitalizations for respiratory infections in multiple hospitals in the city [16]. The first COVID-19 cases were detected in France in late January 2020, with partial lockdowns initiated early in March and a complete national

lockdown by mid-March 2020 [17, 18]. An interrupted time-series analysis based on multicenter prospective surveillance data for pediatric emergency department visits and hospital admissions in Paris demonstrated that the number of visits and admissions after the lockdown decreased by –68% and –45%, respectively. There was also a >70% reduction in acute gastroenteritis, the common cold, bronchiolitis, and acute otitis media compared to the expected values from earlier time periods. However, there was no reduction in visits or admissions associated with urinary tract infections since they were not associated with person-to-person transmission. These data strongly support that social mitigation strategies impact transmission of both contagious respiratory and gastrointestinal pathogens.

Further support for the impact of mitigation strategies was recently published from Hong Kong [19]. This systematic review highlighted the effectiveness of these strategies in reducing influenza burden, largely based on observational studies and simulation exercises. Timely implementation and high compliance in the community are needed to ensure optimal success.

In summary, these studies show that social distancing and mitigation strategies, including closing borders, can make a remarkable difference in infections due to person-to-person transmission. When these measures are no longer indicated for the prevention of SARS-CoV-2 infections, they might be retained in some measure to reduce the burden of other infectious pathogens. However, this will need to be carefully monitored to assess the implications on other aspects of society.

#### Note

**Potential conflicts of interest.** The author has received grants from the Centers for Disease Control and Prevention and the National Institutes of Health; has served as a consultant for Merck, Bio-NET, and IBM; and is on the data and safety monitoring boards of Sanofi, X-4 Pharma, Seqirus, Moderna, and Pfizer. The author has submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant

to the content of the manuscript have been disclosed.

#### References

- Jain S, Williams DJ, Arnold SR, et al; CDC EPIC Study Team. Community-acquired pneumonia requiring hospitalization among U.S. children. *N Engl J Med* 2015; 372:835–45.
- Nair H, Nokes DJ, Gessner BD, et al. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. *Lancet* 2010; 375:1545–55.
- Edwards KM. What have we learned about influenza deaths in children and how can we do better? *Pediatrics* 2018; 141:e20174313.
- World Health Organization. WHO surveillance case definitions for ILI and SARI 2014. Available at: [https://www.who.int/influenza/surveillance\\_monitoring/ili\\_sari\\_surveillance\\_case\\_definition/en/](https://www.who.int/influenza/surveillance_monitoring/ili_sari_surveillance_case_definition/en/). Accessed 7 May 2019.
- Nolen LD, Seeman S, Bruden D, et al. Impact of social distancing and travel restrictions on non-COVID-19 respiratory hospital admissions in young children in rural Alaska [manuscript published online ahead of print 5 September 2020]. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa1328
- Friedrich F, Ongaratto R, Scotta MC, et al. Early impact of social distancing in response to COVID-19 on hospitalizations for acute bronchiolitis in infants in Brazil [manuscript published online ahead of print 28 September 2020]. *Clin Infect Dis* 2021; 72(12):2071–5.
- Yeoh DK, Foley DA, Minney-Smith CA, et al. The impact of COVID-19 public health measures on detections of influenza and respiratory syncytial virus in children during the 2020 Australian winter [manuscript published online ahead of print 28 September 2020]. *Clin Infect Dis* 2021; 72(12):2199–202.
- Foote EM, Singleton RJ, Holman RC, et al. Lower respiratory tract infection hospitalizations among American Indian/Alaska Native children and the general United States child population. *Int J Circumpolar Health* 2015; 74:29256.
- Karron RA, Singleton RJ, Bulkow L, et al. Severe respiratory syncytial virus disease in Alaska native children. RSV Alaska Study Group. *J Infect Dis* 1999; 180:41–9.
- Tumba K, Comaru T, Machado C, Ribeiro M, Pinto LA. Temporal trend of hospitalizations for acute bronchiolitis in infants under one year of age in Brazil between 2008 and 2015. *Rev Paul Pediatr* 2020; 38:7.
- Western Australia Government. COVID-19 coronavirus: WA roadmap [press release]. 2020. Available at: <https://www.wa.gov.au/organisation/department-of-the-premier-and-cabinet/covid-19-coronavirus-wa-roadmap>. Accessed 14 October 2020.
- Agoti CN, Phan MVT, Munywoki PK, et al. Genomic analysis of respiratory syncytial virus infections in households and utility in inferring who infects the infant. *Sci Rep* 2019; 9:10076.
- Kamigaki T, Mimura S, Takahashi Y, Oshitani H. Analysis of influenza transmission in the households of primary and junior high school students during the 2012–13 influenza season in Odate, Japan. *BMC Infect Dis* 2015; 15:282.
- World Health Organization. Influenza update–375. 2020. Available at: [https://www.who.int/influenza/surveillance\\_monitoring/updates/latest\\_update\\_GIP\\_surveillance/en/](https://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/). Accessed 14 October 2020.

15. Olsen SJ, Azziz-Baumgartner E, Budd AP, et al. Decreased influenza activity during the COVID-19 pandemic—United States, Australia, Chile, and South Africa, 2020. *MMWR Morb Mortal Wkly Rep* **2020**; 69:37.
16. Angoulvant F, Ouldali N, Yang DD, et al. COVID-19 pandemic: impact caused by school closure and national lockdown on pediatric visits and admissions for viral and non-viral infections, a time series analysis [manuscript published online ahead of print 3 June 2020]. *Clin Infect Dis* **2021**; 72(2):319-22.
17. Ghanchi A. Adaptation of the national plan for the prevention and fight against pandemic influenza to the 2020 COVID-19 epidemic in France [manuscript published online ahead of print 7 April 2020]. *Disaster Med Public Health Prep* **2020**. doi:10.1017/dmp.2020.82
18. French Government. Décret n° 2020-260 du 16 mars 2020 portant réglementation des déplacements dans le cadre de la lutte contre la propagation du virus COVID-19 [in French]. Available at: <https://www.legifrance.gouv.fr/eli/decret/2020/3/16/PRMX2007858D/jo/texte>. Accessed 14 October 2020.
19. Fong MW, Gao H, Wong JY, et al. Nonpharmaceutical measures for pandemic influenza in nonhealthcare settings—social distancing measures. *Emerg Infect Dis* **2020**; 26:976-84.